


Number 34

Price 20 cents

REPRINT AND CIRCULAR SERIES
OF THE
NATIONAL RESEARCH
COUNCIL



INDEXING OF SCIENTIFIC ARTICLES

BY GORDON S. FULCHER
Corning Glass Works, Corning, New York

Including an Analytic Index
of the *Astrophysical Journal*, Vol. 51-54, 1920-1921

A
0
0
1
2
1
7
3
1
7
5



Announcement Concerning Publications

of the

National Research Council

The Proceedings of the National Academy of Sciences

has been designated as the official organ of the National Research Council for the publication of accounts of research, committee and other reports, and minutes.

Subscription rate for the "Proceedings" is \$5.00 per year. Business address: Home Secretary, National Academy of Sciences, Smithsonian Institution, Washington, D.C.

The Bulletin of the National Research Council

presents contributions from the National Research Council, other than proceedings, for which hitherto no appropriate agencies of publication have existed.

The "Bulletin" is published at irregular intervals. The subscription price, postpaid, is \$5.00 per volume of approximately 500 pages. Numbers of the "Bulletin" are sold separately at prices based upon the cost of manufacture.

The Reprint and Circular Series of the National Research Council

renders available for purchase, at prices dependent upon the cost of manufacture, papers published or printed by or for the National Research Council.

Orders for the "Bulletin" or the "Reprints and Circulars" of the National Research Council, accompanied by remittance, should be addressed: **Publication Office, National Research Council, 1701 Massachusetts Avenue, Washington, D.C.**

INDEXING OF SCIENTIFIC JOURNALS

By GORDON S. FULCHER

The importance of the service which the subject-indexes of journals may render to scientific research is not generally realized. Yet, in order that the best methods may be chosen and unnecessary duplication avoided, each research should, of course, be based on as complete a knowledge of past results as may be obtained; and the task of guiding the scientist to those parts of the enormous accumulation of scientific literature which relate to his work naturally devolves largely upon the subject-indexes. Also, since the greater the amount of time and effort required by scientists to secure the information needed for effective research, the greater will be the proportion discouraged from attempting it and the greater will be the extent to which the research done will be either less efficient or more delayed than it should be, therefore the responsibility resting upon these indexes, particularly the indexes of abstracting and listing journals, to perform this important task well, is very great.

The question then arises, What characteristics must the indexes have if they are to give the maximum service?

The indexes now provided are of various types and degrees of usefulness. The simplest is a mere *unclassified index of titles*, which are arranged alphabetically by their significant words. The indexes of the *Astrophysical Journal*, the *Physical Review*, and many other journals are of this type.

An improved type is the *classified index of titles* in which the titles are arranged under a limited number of subject headings so as to bring related titles together, the classification being based, however, wholly on the titles. The indexes of *Science Abstracts*, *Journal de Physique*, and others are of this type.

The inadequacy of both these types is obvious for it is well known that in most instances a title cannot sufficiently describe all the subjects treated in the article. A paper on "The Atomic Weight of Iodine" may contain results relating to I_2O_5 and to the occlusion of oxygen by glass; and one on "The Flora of Formosa" may describe new species and perhaps new genera.

A much more serviceable type is the *index of titles classified by content of the articles*, which, while retaining the titles, recognizes their shortcomings and classifies each with more or less precision under headings determined by an examination of the article itself or an abstract. Thus "The Flora of Formosa" would be listed under the genera, species, and other subjects concerning which new information is given. The *International Catalogue of Scientific Literature* and the cumulated indexes of the H. W. Wilson Company—*Agricultural Index*, *Industrial Arts Index*, etc.—are of this character. The value of this type of index is enhanced if, as in the case of the *Engineering Index* and the card indexes of the Concilium Bibliographicum, the classified titles are supplemented by brief statements as to the contents.

Since, however, most titles are incomplete and many inaccurate, why not disregard them altogether? This is done by the *content index*. In preparing it, the precise subjects dealt with in each article are determined by an analysis of the original or of a reliable abstract, index entries which adequately describe these contents are formulated, and finally these entries are indexed under the proper subject-headings, alphabetically arranged. The index of *Chemical Abstracts* is of this type, being based on an analysis of the abstracts, and is not only one of the largest but also one of the most generally useful indexes now being issued. For the most part, however, the classification is carried through one stage only, with the result that if information on a certain phase of one of the larger subjects is desired, many references may have to be looked up. Moreover, the index depends on abstracts which vary considerably in their standards.

The final step in the development of the subject-index is to base the entries directly—or indirectly through abstracts—on a thorough analysis of the original articles, and to carry the subclassification of the entries through two or three stages so as to make each subdivision complete. Since the articles in the *Astrophysical Journal* for 1920 and 1921 were carefully analyzed in preparing the analytic abstracts¹ which precede the articles, it was possible to prepare

¹ Gordon S. Fulcher, "Scientific abstracting," *Science*, 54, 291-95, 1921.

from them a very thoroughly classified *analytic index*. This is reprinted at the end of the paper.

To show the characteristics of the analytic index as compared with those of two types of title index, the entries used by each type to index the same four articles, all relating to photography (*Astrophysical Journal*, 52, 86; 52, 98; 52, 201; and 53, 349), are collected here:

Index of the Astrophysical Journal
(*Unclassified Title Index*)

Contraction and Distortion on Photographic Plates, Image
Image Contraction and Distortion on Photographic Plates
Images, Mutual Action of Adjacent Photographic
Photographic Images, Mutual Action of Adjacent
Photographic Plates, Image Contraction and Distortion on
Photographic Sharpness and Resolving Power
Photometry and the Purkinje Effect, Photographic
Purkinje effect, Photographic Photometry and the
Resolving Power, Photographic Sharpness and

Index of Science Abstracts
(*Classified Title Index*)

Photography

Images, Mutual Action of Adjacent Photographic¹
Photometry, Photographic, and Purkinje-Effect
Plates, Photographic, Image Contraction and Distortion on
Resolving Power, Photographic Sharpness and¹

Vision

Purkinje-effect, and Photographic Photometry

*Analytic Index*²

Photographic plates; properties

contrast functions

drying, time of

grain size

images

contraction and distortion

mutual action

sections

¹ These references are put in by analogy with the others, as the 1921 index has not yet been issued.

² Headings and subheadings alone are given; see the complete index herewith reprinted for details.

- intensification
- penetration of light
- resolving power
- sharpness
- shifts of spectrum lines and star images
- theoretical relations
- turbidity
- Photometry*, photographic, heterochromatic
- Purkinje effect*, photographic
- Spectra*
 - measurements
 - errors possible
 - shifts of lines
 - photographic
- Stars*
 - measurements from photographs
 - errors possible
- Sun*
 - spectrum
 - shifts of lines
 - photographic

The precision and completeness of the analytic index as compared with the title indexes is evident. Moreover, since all the results relating to each subject are indexed together, the analytic index also provides a summary, in outline, of the work done in each field; that is, the reader interested in a particular subject, such as clusters, nebulae, spectra, variables, can determine in what phases of the subject progress has been made by consulting the proper sections of the index. On the other hand, the danger of indexing by titles alone is illustrated by the classification, in the *Index of Science Abstracts*, of the reference to the photographic Purkinje effect under "Vision" whereas the article contains, in fact, no reference to the eye, the indexer being misled by the title.

It may be objected that, while the analytic and other good content indexes are clearly more useful than any title indexes, they are too long and too difficult and laborious to prepare. But a comparison of the analytic index, here reprinted, with the corresponding four separate volume indexes of the *Astrophysical Journal* shows that the analytic index has only twice as many words as the

unclassified title indexes, though containing three times as many references, which are, moreover, precisely and thoroughly classified. Also, the preparation does not require remarkable ability nor should it take an undue amount of the editor's time if the work is well organized. After deciding as to the classification to be adopted, the editor's task is merely to indicate on the abstracts the titles to be used to index the articles (heads and subheads being designated in some convenient way) and later to go carefully through the entries which have been typewritten on slips and arranged alphabetically. The clerical work can be greatly simplified by the use of methods developed by such indexing agencies as *Chemical Abstracts*, the H. W. Wilson Company, and the *New York Times*.

The subject-indexes now serving the various sciences vary widely in type and usefulness, as stated above. Some are very poor, some very good; but none is as complete and precise as it should be if it is to give maximum service to research in its field. It is only a question of time, however, when the workers in each science will come to realize the incalculable value of an efficient abstract journal combined with a complete, precise, and thoroughly classified subject-index, and they will arrange to make the small sacrifices of time and money necessary to secure such uniquely important tools.

ANALYTIC INDEX OF *ASTROPHYSICAL JOURNAL* FOR 1920 AND 1921, VOLUMES 51-54, BASED ON THE ANALYTIC ABSTRACTS

For references to spectra and spectroscopy of elements and compounds *see* Spectra.

Absorption of light

by atmosphere, 315-290 $\mu\mu$; coefficients, 54, 297
by vapors in King's electric furnace; Kirchhoff's law applied to, 51, 13

Aethylamine, preparation of; method, 52, 129

Angles, extremely small; interferometer method of measurement, 51, 257

Arc

low-current; anode fall, cathode fall, potential and potential gradient as functions of current, to 1 amp., for various cathode materials (Ag, C, Cd, Cu, Fe, Sn, Zn, and salts of Ba, Ca, Cs, Na, and Sr), 54, 273

Atmosphere (*see also* Sky)

absorption of light, 315-290 $\mu\mu$; coefficients, 54, 297
absorption spectrum (Fraunhofer lines) (*see* Sun, spectrum)
dispersion of light; measurement; stellar interferometer method suggested, 51, 263
ozone in; amount, daily variations, location, and suggested origin, 54, 297
Atomic theory, Bohr (*see* Spectra, theory)

Benzene; dispersion, rotary magnetic, 436-620 $\mu\mu$; 54, 45

- Benzene, nitro-**; dispersion, rotary magnetic, 436-620 μ , 54, 45
- Binaries, spectroscopic** (*see also* Variables)
 Boss 373, elements of both orbits, parallax, and proper motion, 53, 201
 Boss 3644, Virginis, elements of orbit, 54, 226
 Boss 5026, elements of both orbits, 51, 187
 Boss 5591, elements of both orbits, parallax, and proper motion, 53, 201
 +65°369 Camelopard, radial velocity, 52, 198
 α Capella, elements determined by interferometer, 51, 263
 49 δ Capricorni, elements of orbit, 54, 127
 12, 15K, 42, and 50 Cassiopeiae, radial velocities, 52, 198
 19 Cephei, radial velocity variations, 51, 252
 13 Ceti, binary component; elements of orbit, 52, 110
 Class Oe5; 19 Cephei, A Cygni, 9 Sagittae, and 4 more; radial velocity variations, 51, 252
 A Cygni, radial velocity variations, 51, 252
 X Cygni (Cepheid), spectroscopic orbit, 53, 95
 65 τ Cygni, binary component, elements of orbit, 53, 144
 34 ψ , 68, 71, +62°1637 Draconis, radial velocities, 52, 198
 205 Draconis, elements of both orbits, parallax and proper motion, 53, 201
 dynamics; tides on sphere due to second sphere rotating around it, 51, 309
 comp. α Hercules, elements of orbit, parallax and proper motion, 53, 201
 Lalande 29330 and 46867, elements of orbit, parallax and proper motion, 53, 201
 magnitudes, absolute (*see* parallax)
 measurement of relative brightness and position of components; interferometer method, 51, 257, 263
 orbits; elements of, including periods, radial velocity curves, mass functions (*see* Boss 373, 3644, 5026, 5591; α Capella, 49 δ Capricorni, 13 Ceti; X and 65 τ Cygni, 205 Draconis, comp. α Hercules, Lalande 29330 and 46867, O Σ 82)
 measurement of; interferometer method, 51, 257
 origin, theory of, from nebulae, 51, 309
 π 4 Orionis, photometric study, 51, 218
 O Σ 82, elements of orbit, parallax, and proper motion, 53, 201
 parallax and absolute magnitude (*see* Boss 373, 5591; 205 Draconis, comp. α Hercules, Lalande 29330 and 46867, O Σ 82, 53, 201)
 proper motion (*see*: Boss 373, 5591; 205 Draconis, comp. α Hercules, Lalande 29330 and 46867, O Σ 82, 53, 201)
 radial velocity (*see* orbits, *also* +65°369, Camelopard; 12, 15K, 42, and 50 Cassiopeiae, Class Oe5; 34 ψ , 68, 71, +62°1637 Draconis, 21 π Ursae Minoris)
- radiation pressure between components, theory, 53, 1
 9 Sagittae, radial velocity variations, 51, 252
 spectrum of Boss 2830, comp. α Geminorum, and W Serpentis; notes, 53, 13
 theory (*see* dynamics, origin, and radiation pressure)
 21 π Ursae Minoris, radial velocity, 52, 198
- Binaries, visual (Double stars)**
 13 Ceti, orbit, 52, 110 (*see* Triple systems)
 α Hercules and comp.; probably optical pair, 53, 201
 measurement of separation, photographic; possible errors due to mutual action, 53, 349
- Brightness** (*see* Clusters, Comet, Galaxy, Magnitudes, Nebulae, Novae, and Sky)
- Brightness ratio**, of palladium point to gold point, 51, 244
- Camphor**
 dispersion, rotary; magnetic and natural; solution in ethyl alcohol, 54, 116
 refractive indices for 1:1 solution in ethyl alcohol; 436-620 μ , 54, 116
- Carbon disulfide**; dispersion, rotary, magnetic; 436-620 μ , 54, 45
- Cathode**, for vacuum tube
 carbon, limed; preparation, 53, 323
 use as source of large current, 53, 323
- Cathode rays**; excitation of light in air; intensity for 1500-3500 volts, 52, 278
- Cepheids** (*see* Variables)
- Cluster stars**
 catalogue, photometric (colors and magnitudes)
 in Messier 3, 848 stars, 51, 140
 in Messier 68, 56 giants, 51, 49
 colors (*see* catalogue)
 distribution (number) (*see* M₃ and M68)
 relation to magnitude (*see* M₃ and M68)
 variation with radial distance (*see* M₃)
 giants (*see* M68)
 color; relation to magnitude, general conclusions, 51, 49
 magnitudes (*see* catalogue)
 distribution (number) (*see* M₃ and M68)
 relation to color (*see* M₃ and M68)
 variation with radial distance (*see* M₃)
 in Messier 3; photometric analysis, 51, 140:
 colors and magnitudes
 catalogue of 848 stars
 distribution (number of each color and magnitude)
 distribution in space, variation with distance from center
 relation between color and magnitude
 position co-ordinates of 370 stars
 variables, 17 probable
 in Messier 53 (N.G.C. 5024); variables
 positions of 23 shown in photograph, 52, 73
 in Messier 56 (N.G.C. 6779); variables
 positions of 3 shown in photograph, 52, 73

Cluster stars (*cont.*)

- in Messier 68 (N.G.C. 4590); photometric analysis, **51**, 49;
 - giants, colors, and magnitudes catalogue of 56
 - distribution (number of each color and magnitude)
 - relation between color and magnitude
 - variables; colors, magnitudes, positions, and ranges of variation of 28
 - in Messier 72 (N.G.C. 6891)
 - comparison stars, magnitudes of 29, **52**, 232
 - variables; periods and light curves of 26
 - Cepheids, **52**, 232
 - position of 34, in photograph, **52**, 73
 - in Messier 75 (N.G.C. 6864); variables
 - position of 16, in photograph, **52**, 73
 - in N.G.C. 7006 and 7789; number, **52**, 73
 - number, in N.G.C. 7006 and 7789, **52**, 73
 - number of each color and magnitude (*see* M3, M68)
 - variables (*see* M3, 53, 56, 68, 72, 75; Variables)
- Clusters** (*see also* Cluster stars)
- analysis, photometric (colors and magnitudes of the stars) (*see* Messier 3 and 68)
 - brightness of average cluster, **52**, 73
 - diameter, of N.G.C. 7006 and 7789, **52**, 73
 - form of Messier 68, **71**, 49
 - magnitude, absolute (*see* parallax)
 - Messier 3; photometric analysis, **51**, 140
 - Messier 8 (N.G.C. 6530), photograph, **51**, 4
 - Messier 16 (N.G.C. 6611), photograph, **51**, 4
 - Messier 22 (N.G.C. 6656), photograph, **51**, 4
 - Messier 53 (N.G.C. 5024), photograph, **52**, 73
 - Messier 56 (N.G.C. 6779), photograph, **52**, 73
 - Messier 68 (N.G.C. 4590); analysis, photometric; form; computed parallax; photograph, **51**, 49
 - Messier 72 (N.G.C. 6891); parallax, **52**, 232; photograph, **52**, 73
 - Messier 75 (N.G.C. 6864); photograph, **52**, 73
 - N.G.C. 7006; diameter, parallax and star counts, **52**, 73
 - N.G.C. 7492, photograph, **52**, 73
 - N.G.C. 7789; diameter; number of stars, **52**, 73
 - parallax and absolute magnitude (*see* Messier 68, 72; N.G.C. 7006)
 - average cluster, brightness, **52**, 73
 - computation, methods, **51**, 49
 - globular clusters, forty, **52**, 73
 - photographs, large scale (60-inch)
 - Messier 53, 56, 72, 75 and N.G.C. 7492, showing variable stars, **52**, 73
 - Messier 68, showing variables, **51**, 49
 - in Sagittarius and Scutum, including Messie 8, 16, and 22, **51**, 4
 - stars in (*see* Cluster stars)

Colors (*see* Cluster stars, Novae, Stars, Variables)

Comets

- 1919b; brightness, photographs, behavior of tail, **51**, 103
- Morehouse's; rejection of tail, **51**, 103
- tail, rejection of; instances and stages, **51**, 103
- Constant stars (*see* Stars)
- Córdoba Observatory Catalogue
 - correction for position of star 12431, **51**, 4
- Corona (*see* Sun)

Dark markings (*see* Nebulae, dark)

Diameters, of planetoids, satellites and stars

- measurement (*see* Interferometer, stellar)

Discharge through gases

occlusion of RaEm, **54**, 285

Dispersion

- by atmosphere; measurement; interferometer method suggested, **51**, 263
- theory, electron, modified to include variation with temperature, **51**, 223; note as to priority, **53**, 326

Dispersion, rotary (*see* Camphor, Limonene, Sugar, Tartaric acid, **54**, 116)

Dispersion, rotary, magnetic (*see* Benzene, nitro-Benzene, Camphor, Carbon disulfide, Ethyl iodide, Limonene, α -monobrom-Naphthalene, Sugar, Tartaric acid)

- theory, electron, extended
 - isotropic transparent media, **54**, 45
 - optically active media, **54**, 116

Double stars (*see* Binaries, visual)

Dynamics

- binary system (*see* Binaries, spectroscopic)
- single mass, rotating; equilibrium, **51**, 309
- potential of distorted ellipsoid, **51**, 309

Eclipse (*see* Sun)

Einstein effect (*see* Gravitation)

Electron theory (*see* Dispersion and Dispersion, rotary, magnetic)

Electrons; e/m for those active in magnetic rotary dispersion. computed, **54**, 45

Emission of light, in discharge tube

intensity as function of energy of exciting cathode rays, **52**, 278

Equilibrium (*see* Dynamics)

Ethyl iodide; dispersion, magnetic rotary, 436–620 $\mu\mu$, **54**, 45

Exploded wire, source of high temperature spectra (*see* Spectra)

appearance and mechanical effects (Plate), **51**, 37

Fluorine; preparation of gas by electrolysis, **54**, 133

Furnace

- black body, double platinum wound, **51**, 244
- electric (*see* Spectra)
- vacuum, cathode ray, for high temperature, **53**, 323

Galaxy

- brightness, surface, as viewed from distance, **52**, 162

Galaxy (*cont.*)

comparison with typical spiral nebulae, **52**, 162
 distribution of stars in; section, **52**, 23
 origin; theory, **51**, 309
 radiation pressure exerted on nebulae, **53**, 1

Giant stars (*see* Cluster stars and Variables)**Gratings** (*see* Spectra)

polarizing effect on reflected and transmitted light, **51**, 129
 theory, Rayleigh-Voigt; evidence confirming, **51**, 129

Gravitation

attraction on nebulae due to Galaxy, **53**, 1
 Einstein effect; photographic measurement; possible error due to mutual action of adjacent images, **52**, 98; **53**, 349
 Majorana's theory; astronomical consequences, **54**, 334

Hydrogen; production by high potential sparks, apparent, **52**, 47

Integral, definite

table of values of $\int_0^1 (1-x^2)^{n+\frac{1}{2}} \cos kx \cdot dx$, **53**, 249

Interferometer, stellar (Michelson's)

description
 100-inch; accuracy and limits, **51**, 263
 20-foot; construction, adjustments, and photographs, **53**, 249
 modification to compensate for atmospheric dispersion, **51**, 263; **54**, 78
 effective wave-length, determination, **51**, 263
 measurements
 α Capella; separation and positions of components, and inclination of orbit, **51**, 263
 theory and uses; to measure
 angles, extremely small, and changes of angle, **51**, 257
 diameters of planetoids, satellites, and stars, **51**, 257; *also* distribution of luminosity on the disks, **53**, 249
 dispersion of the atmosphere, **51**, 263
 parallaxes and relative motions of stars, **51**, 257
 relative brightness, position, and separation of double stars (binaries), **51**, 257, 263

Jean's contributions to theory of cosmic origins, **51**, 309

Kirchhoff's law (*see* Absorption)

Light (*see* Absorption, Dispersion, Emission, Polarization, Radiation, Refraction, Spectra)

Limonene

dispersion, rotary, magnetic, and natural, 436–620 μ , **54**, 116
 refractive indices, 436–620 μ , **54**, 116

Magnetic rotary dispersion (*see* Dispersion)

Magnitudes (*see* Binaries, Clusters, Cluster stars, Stars, Variables)

Markings, dark (*see* Nebulae)

Mass, constancy of; possible influence of one body on the mass of another, **54**, 334

Melting-point, of palladium, **51**, 244

Meteorites, falling into sun; light to be expected from, **51**, 37

Molecular models, Bohr type, of halogen acids, as basis for theory of band structure, **51**, 230

Naphthalene, α-monobrom-; dispersion, rotary, magnetic, 436–620 μ , **54**, 45

N.G.C.; corrections to descriptions and positions of various nebulae, **51**, 276

Nebulae

attraction, gravitational, by Galaxy, **53**, 1
 brightness, surface, of nebulous areas
 measurement, photographic method, **52**, 162
 spirals; results for well-known nebulae compared with value for Galaxy, **52**, 162
 catalogue, descriptive, of 330, **51**, 276
 changes in N.G.C. 1555 and 2245, **51**, 276
 constant (unchanged); N.G.C. 995, 1186, 2024, and 7023, **51**, 276
 dark clouds or markings; photographs
 I.C. II, 5146, Cygnus; unique array, **51**, 276
 N.G.C. 2146, Camelopard, **51**, 276
 near γ Orionis, including Barnard **33**, **53**, 392
 in Sagittarius and Scutum; N.G.C. 6523 (M8), 6611 (M16), and 6618 (M17), **51**, 4
 forces acting on; electrostatic, gravitational, and radiative, **53**, 1
 internal motion in spirals (*see* Messier **51**, 81)
 I.C. 431, 432, and 434; photographs, **53**, 392
 I.C. II, 5146, Cygnus; dark markings; photograph, **51**, 276
 Messier 8 (N.G.C. 6523); photograph, **51**, 4
 Messier 16 (N.G.C. 6611); photograph and spectrum, **51**, 4
 Messier 17 (N.G.C. 6618) (Swan); photograph, **51**, 4
 Messier **51** (spiral); internal motion, photograph, and proper motion, **54**, 237
 Messier 81 (spiral); internal motion, photograph, and proper motion, **54**, 347
 N.G.C. 1555 and 2245; changes in, **51**, 276
 N.G.C. 1700 and 3379; radial velocity and spectrum, **51**, 276
 N.G.C. 2023 and 2024; photograph, **53**, 392
 N.G.C. 2146; photograph, **51**, 276
 N.G.C. 2261 (Hubble), theoretical explanation, **53**, 169
 N.G.C.; corrections to descriptions and positions of various nebulae, **51**, 276
 new; descriptive catalogue of 255, **51**, 276
 origin, theory of
 N.G.C. 2261, with fan-shaped appendage, **53**, 169

Nebulae (*cont.*)

- origin, theory of (*cont.*)
 - planetary nebulae, by collision, **54**, 229
 - spirals, **54**, 347; Jeans' theory, **51**, 309
- photographs, large scale (*see* I.C. 431, 432, 434;
I.C. II, 5146; M8, 16, 17, **51**, 81; N.G.C.
2023, 2024, 2146; *also* dark clouds)
- near ζ Orionis, **53**, 392
- in Sagittarius and Scutum, **51**, 4
- spirals; N.G.C. 2146 and 15 others, **51**, 276;
M51, **54**, 237; M81, **54**, 347
- misc.: I.C. I, 1470; N.G.C. 1491, 2245, 2247,
2294, 2359, 3379, 3384, and 6888, **51**, 276
- planetary, origin; collision theory, **54**, 229
- proper motion (*see* Messier 51 and 81)
- radial velocity
 - explanations suggested; discussion, **53**, 1
 - N.G.C. 1700 and 3379, **51**, 276
 - radiation pressure on, due to Galaxy, **53**, 1
 - spectrum (*see* M16; N.G.C. 1700 and 3379)
 - nebulousity around six nebulous stars, **52**, 8
 - spirals (*see* brightness, catalogue, internal
motion, new, origin, photographs, proper
motion)
 - Swan (N.G.C. 6618); photograph, **51**, 4
 - theory (*see* N.G.C. 2261, origin, radial velocity)
 - effect of passage of a star near or through a
nebula, **53**, 169; **54**, 229
 - types, four; examples, **53**, 392
 - variable (*see* changes)

Nebulous areas; brightness (*see* Nebulae)

Nebulous stars (*see* Stars)

Novae

- brightness and color of Persei No. 2, **52**, 183
- definition suggested, **54**, 229
- Ophiuchi 1919; spectrum and radial velocity,
51, 121
- origin, collision theory, **54**, 229
- spectrum
 - Ophiuchi 1919; shifts of lines and bands,
51, 121
 - shifts, simple interpretation, **51**, 121
- theory (*see* origin)

- Occlusion**, of RaEm in discharge tubes, **54**, 285
- Origin**; theory of (*see* Binaries, Galaxy, Nebulae,
Novae, Solar system, Variables)
- cosmic origins; Jeans' contributions, **51**, 309
- Ozone**, in atmosphere (*see* Atmosphere)

Palladium, melting-point, **51**, 244

Parallax (*see* Binaries, Clusters, Stars, Variables)

Photo-electric cells

- alkali metals and hydrides; color sensitiveness
curves, **52**, 129
- fatigue tests for K and KH cells, **52**, 129
- preparation of Li cell, **52**, 129
- sensitiveness, color, for 30 cells, including all
alkali metals and hydrides of Na, K, Rb,
and Cs, **52**, 129

Photo-electric photometer, for stars

- description, **51**, 193; precision, **53**, 105; **54**, 81

Photographic plates; properties

- contrast functions γ and Γ ; variation with
wave-length; Purkinje effect, **52**, 86

- drying, time of; effect of exposure on, **53**, 349

- grain size; relation to sharpness, theoretical,
52, 201

images

- contraction and distortion during drying,
52, 98

- mutual action of adjacent spectrum lines and
stars; analysis of effects involved; varia-

- tion with exposure, **53**, 349

- sections of star images, Plates, **52**, 86, 98

- intensification; effect on resolving power, **52**, 201

- penetration of light of different wave-lengths,
52, 86

resolving power

- effect of intensification, **52**, 201

- relation to grain size and sharpness, theo-

- retical, **52**, 201

- variation with development and wave length
52, 201

sharpness

- measurement; improved method, **52**, 201

- relation to contrast and turbidity, **52**, 201

- variation with development and wave-length
52, 201

shifts of spectrum lines and star images due to

- mutual action of adjacent images (*see*

- images)

theoretical relations for resolving power and

- sharpness, **52**, 201

- turbidity; variation with wave-length, **52**, 201

Photographs (*see* Clusters, Comets, Nebulae, Sun)

Photometer (*see* Photo-electric photometer)

Photometry, photographic, heterochromatic; dis-

- cussion and warning, **52**, 86

Planetoids, diameter; measurement, interferom-

- eter method, **51**, 257

Polarization

- measurement, using gratings; warning, **51**, 129

- produced by gratings, reflected and transmitted

- light, **51**, 129

Prominences, solar (*see* Sun)

Proper motion (*see* Binaries, Nebulae, Stars)

Purkinje effect, photographic, **52**, 86

Quantum theory (*see* Spectra)

Radial velocity (*see* Binaries, Nebulae, Novae,

- Stars)

Radiation

- constant c_2 ; computed from gold point to

- palladium point ratio, **51**, 244

pressure

- on atoms and electrons; theory based on

- classical dynamics, **52**, 65

- between binary stars, **53**, 1

- on nebulae due to Galaxy, **53**, 1

Radium emanation (Niton)

occlusion in discharge tube, **54**, 285
purification; modification of Duane's apparatus, **54**, 285

Refractive indices (*see* Camphor, Limonene, Sugar, Tartaric acid)**Satellites**, diameter; measurement; interferometer method, **51**, 257**Scandium**, carbide; possible formation in electric furnace, **54**, 28**Sky, night**; brightness; various determinations; discussion, **52**, 123**Solar corona and prominences** (*see* Sun)**Solar system**, origin; tidal theory, **51**, 309**Spectra and spectroscopy**

absorption spectra

exploded wire, of Fe, λ 2270-5645 Å, spectrogram, **51**, 37

furnace, electric, of Ba, Ca, Co, Fe, Ni, and Ti; variation of relative intensities with temperature, **51**, 13

production of

electric furnace spectra, **51**, 13

high temperature spectra, extreme; exploded wire method, **51**, 37

theory (*see* halogen acid gases)

Kirchhoff's law applied to electric furnace spectra, **51**, 13

Zeeman effect, inverse (*see* Fe and V)

air

arc spectrum; two new lines, **54**, 246

spark spectrum, condensed

effect of self-inductance on relative intensities, 590-872 $\mu\mu$, **51**, 236

identification of Ar, N, and O lines, 590-872 $\mu\mu$, **51**, 236; **54**, 76

shift with reference to vacuum tube lines, **51**, 236

wave-lengths, 590-872 $\mu\mu$, **51**, 236; **54**, 76

ammonia bands, visible and ultra-violet; identification, **52**, 301

arc spectra (*see* air, Fe, pole-effect, pressure shift)

anode and cathode spectra; relative behavior of various metal lines; variation with atomic weight, **54**, 65

comparison with furnace spectra (*see* Ca, cyanogen, and Swan bands)

ionization lines; behavior, **54**, 191, 246

low-current; variation with current; excitation stages

for Ba, Ca, K, Mg, Na, Sr, **54**, 191

for Ag, Bi, Cd, Cu, and Zn, **54**, 246

relation of results to Bohr theory and Ritz equations, **54**, 246

standard lines, secondary; comparison of 12-mm 5-amp. with 6-mm 6-amp. arc, **53**, 260

argon; lines in condensed spark spectrum in air, 590-872 $\mu\mu$, **51**, 236; **54**, 76; shift with reference to vacuum tube lines, **51**, 236

band spectra (*see* ammonia, cyanogen, halogen acids, Swan, water, C, CO, CO₂, CF₄, H₂S, N, SO₂)

arc and furnace spectra; comparison of intensity distribution; CN and Swan bands, **53**, 161

Deslandres' law; test with nitrogen positive bands, **52**, 301

theory of structure of infra-red bands (*see* halogen acids)

vacuum tube discharge through CO₂, H₂S, NH₄, N₂O, N₂O₄, and SO₂; bands excited in visible and ultra-violet, **52**, 301

barium

absorption spectrum, electric furnace; variation with temperature, **51**, 13

arc spectrum, low current; variation with current, **54**, 191

classification of lines

furnace excitation, infra-red, **51**, 179

low-current arc excitation, **54**, 191

infra-red furnace spectrum to 856 $\mu\mu$, at various temperatures, **51**, 179

series of single lines and triplets; identification of terms; constants, **51**, 23

binaries (*see* Binaries)

bismuth

arc spectrum, low current; variation with current, **54**, 246

classification of lines, arc excitation, **54**, 246

structure

of $\lambda\lambda$ 4122, 4308, 4722; Plate, **53**, 323, 339

of $\lambda\lambda$ 3397, 3511, 3506; Plate only, **53**, 323

of λ 4722; changes in relative intensity of components, **53**, 339

cadmium

arc spectrum, low current; variation with current, **54**, 246

classification of lines, arc excitation, **54**, 246

new resonance line, λ 3779, **54**, 246

calcium

absorption spectrum, electric furnace, **51**, 13

arc spectrum, low current; variation with current, **54**, 191

classification of lines

furnace excitation, infra-red, **51**, 179

low-current arc excitation, **54**, 191

furnace spectra; absorption, emission and mixed, **51**, 13; infra-red, **51**, 179

infra-red furnace spectrum to 733 $\mu\mu$, at various temperatures, **51**, 179

new lines, fifty, **52**, 265

pressure shifts to 1 atm., 315-650 $\mu\mu$, **53**, 224

series of singlets and triplets; identification of terms; constants, **52**, 265

carbon

bands, negative; origin, **52**, 301

band, positive, fourth; wave-lengths, **52**, 301

ultra-violet vacuum spark spectrum; λ 1931-360 Å; Plate, **52**, 47; wave-lengths, **53**, 150

X-ray spectrum, L-series; identification, **52**, 47

Spectra (*cont.*)

carbon dioxide; bands in visible and ultra-violet, including several new, **52**, 301

carbon monoxide; bands in visible and ultra-violet; identification, **52**, 301

carbon tetrafluoride; bands in visible and ultra-violet; wave-lengths, **54**, 133

cathode rays; intensity of N bands excited, as function of energy, **52**, 278

classification of lines

enhanced lines, Fowler's; discussion, **54**, 246

furnace excitation stages, King's electric;

for Ba, Ca, Co, Ni, and Sr; infra-red, **51**, 179

for Mn, λ 2795-6500 Å, **53**, 133

for Sc, λ 3015-6559 Å, **54**, 29

low-current arc excitation stages

for Ag, Bi, Cd, Cu, and Zn, **54**, 246

for Ba, Ca, K, Mg, Na, and Sr, **54**, 191

comparison with furnace classification, **54**, 246

cobalt

absorption spectrum, electric furnace; variation with temperature, **51**, 13

classification of lines; furnace stages; infra-red to 809 μ , **51**, 179

infra-red furnace spectrum to 809 μ , at various temperatures, **51**, 179

copper

arc spectrum, low current; variation with current, **54**, 246

classification of lines; low-current arc stages and comparison with furnace stages, **54**, 246

continuous background obtained with exploded wire source, **51**, 37

cyanogen bands

intensity distribution, in furnace and arc spectra, **53**, 161

λ 3883 in arc and furnace; Plate, **53**, 161

excitation of spectra; minimum voltage; helium spectra, **52**, 1;

variation with current density; helium, **52**, 1

exploded wire spectra (extremely high temperature absorption and emission spectra)

of Cu, Ni, Mn; continuous background, **51**, 37

of Fe; absorption spectrum, λ 2270-5645 Å; Plate, **51**, 37

production, method, **51**, 37

use for study of pressure shift suggested, **51**, 37

fluorescence spectra (*see* mercury, **54**, 149)

fluorine; spark spectrum, visible and ultra-violet, of pure gas; wave-lengths, **54**, 133

Fraunhofer lines (*see* Sun, spectrum)

furnace spectra, electric

absorption spectra of metallic vapors

comparison with emission spectra, **51**, 13

production, method, **51**, 13

theory; Kirchhoff's law applied to, **51**, 13

variation with temperature; spectra of

Ba, Ca, Co, Fe, Ni, and Ti, **51**, 13

Ba, Ca, Co, Ni, Sr; infra-red, **51**, 179

Mn, 280-820 μ , **53**, 133

Sc, 301-656 μ , **54**, 28

comparison with arc spectra

intensity distribution in

cyanogen and Swan bands, **53**, 161

spectra of Ba, Ca, Co, Ni, Sr, **51**, 179; Sc, **54**, 28

comparison with solar spectrum; Sc, **54**, 28

effect of small potential gradient, **52**, 187

infra-red absorption spectra to 920 μ , of

Ba, Ca, Co, Ni, and Sr, **51**, 179

mixed absorption and emission spectra; production, method, **51**, 13

origin of radiation; discussion, **52**, 187

red fringe; explanation, **52**, 187

variation with temperature (*see* furnace absorption spectra)

Zeeman effect for iron lines, **51**, 107

grating spectrograph

comparison with interferometer, **53**, 260

ghosts and reversals; use of, in accurate measurements, **53**, 260

intensity effect on wave-length nil, **53**, 260

polarizing effect on reflected and transmitted light, **51**, 129

ultra-violet, extreme, **52**, 47, 286; **53**, 150

halogen acid gases (HBr, HCl, HF); bands

HCl band 3.7 μ ; wave-lengths, law of spacing,

evidence of satellites, **53**, 300

theory of structure of infra-red bands

isotopic theory of doublets, **52**, 248

quantum theory, based on simple molecular

model of Bohr type, **51**, 230

helium

excitation of various spectra; minimum voltage, **52**, 1

intensity, relative, of series lines and bands in arc spectrum; variation with voltage, **52**, 1

ultra-violet, extreme, spark spectrum; identification of lines, **52**, 47

hydrogen, Balmer series

shift of H α ; condensed spark in air compared with vacuum spectrum, **51**, 236

variations in relative intensity of lines in spectrum of Class Md variable star, **53**, 135

hydrogen bromide, chloride, and fluoride (*see* halogen acid gases)

hydrogen sulfide; spectrum of discharge through **52**, 301

infra-red spectra (*see* air, Ba, Ca, Co, Ni, Sr, halogen acid gases, sun, water)

elimination of scattered light of shorter wave-lengths, **53**, 121

screen for light to 7200 Å, **53**, 121

interferometer spectrograph

comparison with grating, **53**, 260

ghosts and reversals; use of, for accurate measurements, **53**, 260

intensity effect on wave-length nil, **53**, 260

reduction of measurements, method, **53**, 260

ionization lines, in low-current arc; variation with current, **54**, 191

Spectra (cont.)

iron

- absorption spectrum
 - furnace; variation with temperature, **51**, 13
 - exploded wire, λ 2270-5645 Å; Plate, **51**, 37
 - arc lines, λ 3370-6750 Å; wave-lengths of 1026 lines, measured with grating and interferometer, and compared with Bureau of Standards results, **53**, 260
 - intensity effect for arc lines nil, **53**, 260
 - pole-effect in Pfund arc
 - comparison of 12-mm 5-amp. with 6-mm 6-amp. arc, **53**, 260
 - relation to Zeeman effect nil, **53**, 329
 - ultra-violet spark spectrum to 200 Å; Plate, **52**, 47; wave-lengths, **53**, 150
 - variations in relative intensity of lines in spectrum of Class Md variable stars, **53**, 185
 - Zeeman effect
 - furnace lines; direct and inverse effect for 100 lines; comparison with results for spark lines; Plates, **51**, 107
 - relation to pole-effect nil, **53**, 329
 - isotopes, components due to measurement of separation; displaceable slit-method suggested, **53**, 329
 - theory, for case of bands of HBr, HCl, **52**, 248
- ### magnesium
- arc spectrum, low-current; variation with current, **54**, 191
 - classification of lines; low-current arc stages, **54**, 191
- ### manganese
- classification of lines, 280-650 $\mu\mu$; furnace stages, **53**, 133
 - furnace spectrum, 280-820 $\mu\mu$; variation with temperature, **53**, 133
- ### measurements
- errors, possible, due to mutual influence of adjacent photographic images, **52**, 98; **53**, 349
 - of shifts, minute; displaceable slit method, suggested, **53**, 329
 - of ultra-violet wave-lengths, extremely short, **53**, 150
- ### mercury; fluorescence spectrum, excitation of;
- active molecules; relation to exciting spectrum, **54**, 149
- ### mixed absorption and emission spectra; production in electric furnace, **51**, 13
- ### nebulae; spectra (see Nebulae)
- ### neon; low-voltage spectrum of trace of neon in helium, **52**, 1
- ### nickel
- absorption spectrum, electric furnace; variation with temperature, **51**, 13
 - classification of lines, furnace stages; infra-red to 780 $\mu\mu$, **51**, 179
 - infra-red furnace spectrum to 920 $\mu\mu$; at various temperatures, **51**, 179

ultra-violet spark spectrum to 731 Å; Plate, **52**, 47; wave-lengths, **53**, 150

nitrogen; band spectrum

- excitation by vacuum discharge through N_2O and N_2O_3 , **52**, 301
- intensity as function of energy of the exciting cathode rays, **52**, 278
- new ultra-violet positive bands from low-current arc in air, possible, **54**, 246
- structure; divergence from Deslandres' law, **52**, 301
- wave-lengths of third positive, **52**, 301

nitrogen; line spectrum

- identification of lines in condensed spark in air, 590-870 $\mu\mu$, **51**, 236; **54**, 76
- relative intensity; effect of self-inductance with spark, **51**, 236
- shift of spark lines with reference to vacuum tube lines, **51**, 236

nitrogen peroxide; spectrum of discharge through, **52**, 301

nitrous oxide; spectrum of discharge through, **52**, 301

novae; spectrum (see Novae)

oxygen

- identification of lines in spark in air and O_2 , 590-870 $\mu\mu$, **51**, 236; **54**, 76
- shift of spark lines with reference to vacuum tube lines, **51**, 236

pole-effect

- in iron arc, comparison of 6-mm 6-amp. and 12-mm 5-amp. arcs, **53**, 260
- measurement, displaceable slit-method suggested, **53**, 329
- relation to Zeeman effect, for iron, nil, **53**, 329

potassium

- arc spectrum, low-current; variation with current, **54**, 191
- classification of lines, low-current arc stages, **54**, 191

pressure shift

- calcium arc lines, 315-650 $\mu\mu$, **53**, 224
- source for study; exploded wire suggested, **51**, 37

in stellar spectra, Arcturus, Procyon, **53**, 327

quantum theory (see theory)

radium emanation (niton)

- new lines, 398-745 $\mu\mu$, **54**, 285
- relative intensity of lines; variation during discharge, **54**, 285

resonance lines in low-current arc spectra

- intensity variation with current for Ag, Bi, Cd, Cu, and Zn, **54**, 246
- Ba, Ca, K, Mg, Na, and Sr, **54**, 191

Ritz equations (see series)

scandium

- classification of lines, furnace stages, **54**, 28
- furnace spectrum, 301-656 $\mu\mu$; at various temperatures; comparison with arc, solar and sun-spot spectra, **54**, 28

Spectra (*cont.*)

scandium (*cont.*)

Zeeman effect for lines in sun-spot spectrum, 54, 28

series

in Ba spectrum, singlets and triplets, 51, 23

in Ca spectrum, singlets and triplets, 52, 265

notation; explanation, 51, 23

Ritz equations; relation of low-current arc results to, 54, 246

shifts of lines (*see* spark spectra, Sun)

measurement; displaceable slit method suggested, 53, 329

photographic, due to mutual action of adjacent images, 52, 98; 53, 349

silver

arc spectrum, low current; variation with current, 54, 246

classification of lines; low-current arc stages, 54, 246

sodium

arc spectrum, low current; variation with current, 54, 191

classification of lines; low-current arc stages, 54, 191

Zeeman effect for D-lines; explanation of Woltjer's observations, 51, 107

sources of light (*see* arc, exploded wire, spark, vacuum)

exploded wire; appearance and mechanical effects of explosion, 51, 37

spark spectra (*see* air, argon, fluorine, hydrogen, nitrogen, oxygen)

relative intensity; effect of self-inductance on air and O lines, 51, 236

shift with reference to vacuum tube lines; air, Ar, H, N, and O lines, 51, 236

ultra-violet, extreme (*see* C, Fe, He, Ni, Zn, ultra-violet)

spark, condensed, in vacuum; as source for extreme ultra-violet, 52, 286

spectrographs (*see* grating, interferometer)

standards, international secondary; questioned lines; comparison of 6-mm 6-amp. and 12-mm 5-amp. arcs; pole-effect, 53, 260

stellar spectra (*see* Stars)

strontium

arc-spectrum, low-current; variation with current, 54, 191

classification of lines

arc stages, low-current, 54, 191

furnace stages, infra-red, 51, 179

infra-red furnace spectrum to $920\ \mu\mu$; at various temperatures, 51, 179

structure (*see* bands, bismuth)

sulfur dioxide bands

in visible and ultra-violet, including forty new; wave-lengths, 52, 301

sun (*see* Sun)

Swan band

intensity distribution; $5165\ \text{\AA}$; arc and furnace spectra, 53, 161

theory

Bohr; relation of low-current arc results to, 54, 246

relation of minimum voltage results for helium to, 52, 1

isotopic, of separation of doublets of HBr and HCl, 52, 248

quantum, of structure of band spectra of halogens, based on simple molecular model of Bohr type, 51, 230

titanium; absorption spectrum, electric furnace, at various temperatures, 51, 13

ultra-violet spectra (*see* Ca, CO, CO₂, F, Mn, N, NH₄, N₂O, N₂O₂, SO₂, Sun)

extreme, to $200\ \text{\AA}$ (*see* C, He, Fe, Ni, Zn) measurements of wave-lengths, 53, 150

source; condensed vacuum spark, 52, 286

spectrograph, vacuum, 52, 47, 286; 53, 150

screens for region $290\text{--}315\ \mu\mu$, 54, 297

spectrograph, special double, for solar spectrum, $290\text{--}315\ \mu\mu$, 54, 297

vacuum discharge spectra (*see* CO, CO₂, CF₄, N, Ne, NH₄, RaEm, SO₂)

intensity of N bands as function of energy of cathode rays, 52, 278

minimum voltage for excitation of He spectra, 52, 1

spectra of discharge through CO₂, H₂S, NH₄, N₂O, N₂O₂, SO₂, 52, 301

vacuum sources

cathode rays from treated carbon cathode, used to heat anode, 53, 323

condensed spark, 52, 286

vanadium; Zeeman effect, direct and inverse, for 90 furnace lines; comparison with effect for spark lines, 51, 107

Venus; spectrum (*see* Venus)

water vapor; absorption band, $930\text{--}963\ \mu\mu$, 53, 121

X-ray spectra

L-series of carbon; identification, 52, 47

Zeeman effect (*see* Fe, Na, Sc, V)

furnace lines compared with spark lines, of Fe and V, 51, 107

relation to pole-effect for Fe lines nil, 53, 329

zinc

arc spectrum, low current; variation with current, 54, 246

classification of lines, low-current arc stages, 54, 246

ultra-violet spark spectrum to $316\ \text{\AA}$; Plates, 52, 47, 286; wave-lengths, 52, 286

Stars

atmospheres, pressure in; Arcturus and Procyon, 53, 327

binaries (*see* Binaries and Variables)

brightness (*see* parallax)

Stars (*cont.*)

catalogue (*see* Cluster stars)
 in Pleiades; magnitudes and colors of 821 stars in region 2° square, 54, 323
 spectroscopic parallaxes, magnitudes, type, and proper motion of 1646 stars, 53, 13
 Cepheids (*see* Variables)
 Class Bo-B5; statistical study of 180 stars; mean magnitude, parallax and proper motion, 54, 140
 Class O; collision theory of origin, 54, 229
 Class Oe5 (*see* Binaries)
 Class Md (*see* Variables)
 cluster (*see* Cluster stars)
 colors (*see* Cluster stars)
 determination for nebulous stars, 52, 8
 nebulous stars, 47, including ρ Ophiuchi, σ and 22 Scorpii, 52, 8
 in Pleiades; 753 dwarfs, 54, 323
 comparison stars, for Messier 72; photographic magnitudes of 29, 52, 232
 constant stars, from photometric studies
 Bond 624, 53, 317
 τ and σ Cassiopeiae, 54, 81
 Class B: π^4 Orionis, ξ , ϵ , and μ Tauri, 51, 193
 π^4 Orionis, 51, 193, 218
 ξ , ϵ , and μ Tauri, 51, 193
 l and π Persei, 53, 105
 radial velocity constant (*see* radial)
 density of stars
 in Gal. long. $+32^\circ$, lat. -20° , 52, 73
 in Galaxy; distribution, 52, 23
 in space, as function of parallax and magnitude, 52, 23
 diameters
 measurement with stellar interferometer, 51, 257 (*see* Interferometer)
 α Orionis, 53, 249
 disk (*see* diameter)
 darkening of limb (*see* Variables)
 distribution of luminosity; interferometer method of study, 53, 249
 distribution in space (*see* density)
 number of each magnitude per unit volume, luminosity curve, 52, 23
 double stars (*see* Binaries, visible)
 dwarf stars, in Pleiades (*see* catalogue)
 Galaxy (*see* Galaxy)
 giants (*see* Cluster stars, Variables)
 magnitudes, absolute (*see* parallax)
 magnitudes, photo-electric, of β , δ , l , π Persei, 53, 105
 magnitudes, photographic and photo-visual
 catalogue, of 1646 stars, 53, 13
 Class Bo-B5, 180 stars, 54, 140
 comparison stars
 for Messier 72, 29, photographic, 52, 232
 for Nova Persei No. 2, 36, 52, 183
 in Messier 3 and 68 (*see* Cluster stars)
 in Orion, eight; photo-visual, 53, 317

in Pleiades; statistical study of 821, 54, 323
 scales, photo-visual; comparison of Barnard's with Mount Wilson, 54, 323
 measurements from photographs; possible errors due to contraction effect, 52, 98; and mutual action of images, 53, 349
 nebulous stars
 colors of 47, including ρ Ophiuchi, σ and 22 Scorpii, 52, 8
 spectrum
 R Aquarii, nebulous lines; Plate, 53, 375
 nebulousity around six stars, 52, 8
 novae (*see* Novae)
 number of each absolute magnitude; luminosity curve, 52, 23 (*see* density)
 parallaxes and absolute magnitudes
 catalogue for 1646 stars; spectroscopic and trigonometric results, 53, 13
 Class Bo-B5 stars; mean for 180, 54, 140
 Boss 1517, 51, 254
 determination; methods
 accuracy, relative; discussion, 54, 140
 interferometer method suggested, 51, 257
 spectroscopic; description, 53, 13
 distribution of stars in space and number of each magnitude, 52, 23
 relation of magnitude to space velocity; statistical study of 1350 stars, 54, 9
 relation of parallax to apparent magnitude and proper motion, 52, 23
 pressure in atmospheres (*see* atmospheres)
 proper motion
 catalogue, for 1646 stars, 53, 13
 Class Bo-B5, mean for 180 stars, 54, 140
 large; two faint stars near M51, 54, 237
 measurement; interferometer method suggested, 51, 257
 radial velocity (*see* velocity, space)
 binaries (*see* Binaries)
 16, 19, $+59^\circ 2395$, $+83^\circ 104$ Cephei, 52, 198
 constant for δ Ophiuchi, η Serpentis, β Pegasi, 52, 317
 $+73^\circ 835$ Draconis, 52, 198
 α Hercules and comp., 53, 201
 25 θ Ursae Majoris, 23 δ Ursae Minoris, 52, 198
 spectra (*see* nebulous stars)
 combined bright and dark line spectra; explanation, 51, 13
 variable (*see* Variables)
 spectral type
 catalogue for 1646 stars, 53, 13
 theory; effect of passage of a star near or through a nebula, 53, 169; 54, 229
 origin of Class O stars, 54, 229
 variable stars (*see* Variables)
 velocity; radial, tangential and space
 distribution; frequency of each velocity; statistical study of 1350 stars, 54, 9
 relation to absolute magnitude, 54, 9

Sugar, cane; aqueous solution
dispersion, rotary, magnetic and natural, 436-620 μ , 54, 116
refractive index, one to one solution, 436-620 μ , 54, 116

Sun

corona; May 29, 1919; photograph, 51, 1
eclipse; May 29, 1919; Smithsonian expedition; brief report, 51, 1 (*see* corona)
meteorites falling into; light expected from; theoretical discussion, 51, 37
prominence; May 29, 1919; photograph, 51, 1
October 8, 1920; very high; stages of growth; photographs with Ca line, 53, 310
spectrum, including Fraunhofer lines
comparison with arc and furnace spectra; scandium lines, 54, 28
energy distribution, corrected for atmospheric absorption, 315-290 μ , 54, 297
infra-red, 890-990 μ ; identification and origin of lines; wave-lengths of 563 lines, including 50 solar; also Plate, 53, 121
shifts of lines
atmospheric refraction, 53, 380
center-arc, predicted by anomalous dispersion theory, 54, 92
mutual influence of adjacent lines; data from limb-center comparisons, 54, 92 (*see* photographic)
photographic effect of adjacent lines, 52, 98; 53, 349
theory, anomalous dispersion; of gravitational shift, limb-center shift, and mutual influence shift, 54, 92
spot spectra (*see* spots)
theory (*see* shifts)
ultra-violet (*see* energy distribution)
map, photographic; 315-290 μ , 54, 297
spots
spectrum
comparison with arc and furnace spectra; scandium lines, 54, 28
Zeeman effect for scandium lines, 54, 28
theory; cooling of rising gases, 54, 293
theory (*see* spectrum, spots)

Tartaric acid

aqueous solution
dispersion, rotary, natural, and magnetic, 436-620 μ , 54, 116
refractive index, for one to one solution, 436-620 μ , 54, 116

Telescope objective; diffraction by; effect on image of disk and combination of disks, including lune; mathematical theory, 51, 73

Temperature scale

brightness ratio, gold point to palladium point, 51, 244
palladium melting-point, 51, 244

Theory (*see* Binaries, Dynamics, Electron, Gratings, Gravitation, Interferometer, Nebulae, Novae, Origin, Photographic plates, Radiation pressure, Spectra, Stars, Sun, Variables

Transits; observation; effect of diffraction by telescope objective; theory, 51, 73

Triple systems

65 τ Cygni; orbit of spectroscopic binary component, 53, 144
13 Ceti; orbit of spectroscopic binary component and perturbations due to fainter visual component, 52, 110
light curve of variable binary component of λ Tauri, 51, 193
orbits; of binary components (*see* 65 τ Cygni, 13 Ceti, and λ Tauri)
perturbations due to third body (*see* 13 Ceti and λ Tauri)
 λ Tauri; photometric study of variable binary component; orbit, light curve, effect of third body nil, 51, 193

Variable nebulae (*see* Nebulae)

Variable stars

Algol; photometric study, elements of eclipsing system, light curve, color of satellite, 53, 105
R Aquarii; spectrum, intensities and displacements of lines and nebular lines (Plate), 53, 375
 γ Argus; spectrum; temporary, shifting, absorption He lines, 52, 39
 γ Camelopardalis; photometric study; light curve and elements, 54, 217
RS Canum Venaticorum; light curve, elements, computed parallax, 53, 99
l Carinae; periodic variations of wave-length and spectral type, 54, 161
x H. Cassiopeiae; photometric study; light curve, elements, darkening of limb, 54, 81
SX Cassiopeiae; light curve, elements, 53, 165
T Cephei; periodic spectrum changes, 53, 185
U Cephei; photometric study; light curve, elements, evidence of tidal evolution, 52, 145
Cepheids (*see* l Carinae, X Cygni, Messier 72)
light curves of 26, in Messier 72, 52, 232
light range, small; possibility of Cepheids with; suggestion, 51, 62
orbit, spectroscopic, for X Cygni, 53, 95
origin; collision theory of, 54, 229
magnitude, mean, for 26 in M72, 52, 232
periods; in M72, 26 variables, 52, 232
relation to spectral type, 54, 161
spectral type; range of variation and relation to period, 54, 16
spectrum; periodic variations in wave-length and type; l Carinae, 54, 161
theory (*see* origin)
general conclusions; mean atomic weight; ratio of mass to radius, 52, 73
relation of period to brightness, 52, 73
variation
binary theory; discussion, 51, 62
condition of; ratio mass to radius, 52, 73

Variable stars (*cont.*)

Class Md; spectrum; periodic changes in emission lines (see T Cephei, X Cygni, R Hydrae, R Leonis, X Ophiuchi, R Serpents, 53, 185; R Aquarii, 53, 375)
classification of long-period variables, 53, 179
cluster variables (*see* Cepheids)
color variation of typical, 51, 49
colors and magnitudes of 28 in M68, 51, 49
magnitude, absolute, of typical, 51, 49
new; in M3; 17 probable, 51, 140
in M53, 56, 72, and 75; positions of 80 shown on photographs, 52, 73
in M68; 28, mostly typical, 51, 49
colors (*see* cluster)
satellite of Algol, 53, 105
X Cygni; spectroscopic orbit, 53, 95
X Cygni; periodic changes of spectrum, 53, 185
darkening of limb; γ H. Cassiopeiae, 54, 81
205 Draconis, probable eclipsing variable, 53, 201 (*see* Binaries)
eclipsing variables
light curve and elements (*see* Algol, γ Camelopardalis, RS Canum Venaticorum, SX and γ H. Cassiopeiae, U Cephei, RT Lacertae, λ Tauri)
probable, 205 Draconis, 53, 201
ellipsoidal variable (*see* π s Orionis)
evolution, tidal, of U Cephei; evidence, 52, 145
giants (*see* Cepheids)
R Hydrae; variations in spectrum, 53, 185
irregular variables; collision theory of origin, 54, 229
RT Lacertae; light curve and elements, 52, 257
R Leonis; variations in spectrum, 53, 185
light curves (*see* Cepheids, eclipsing, ellipsoidal, and long-period variables)
long-period variables
classification, 53, 169
light curves of 66; constants, 53, 169
spectrum of R Aquarii, 53, 375
magnitude (*see* Cepheids)

magnitude, absolute (*see* cluster, parallax)
Messier 3, 53, 56, 68, 72, 75; new variables (*see* cluster)
Messier 72; light curves of 26 Cepheids, 52, 232
nebulous variable (*see* R Aquarii)
new variables (*see* cluster variables)
B.D. $+81^{\circ}27$; $+81^{\circ}30$, 52, 145
orbits; elements (*see* eclipsing variables)
spectroscopic; X Cygni, 53, 95
origin, of Cepheids and irregular variables;
collision theory, 54, 229
X Ophiuchi; variations in spectrum, 53, 185
 π s Orionis; photometric study; light curve and elements, 51, 218
parallax and absolute magnitude, of RS Canum Venaticorum, 53, 99
periods (*see* light curves, Cepheids)
photometric study (*see* Algol, γ Camelopard, γ H. Cassiopeiae, U Cephei, λ Tauri)
 π * Orionis, constant star, 51, 103, 218
R Serpents; variations in spectrum, 53, 185
spectrum; periodic variations in emission lines (*see* R Aquarii, γ Argus, Cepheids, Class Md)
suspected variables
B.D. $+10^{\circ}1771$, 52, 9
Bond 624 in Orion; photometric study;
Hartwig's elements incorrect, 53, 317
205 Draconis, 53, 201
 δ Persei, 53, 105
 λ Tauri; photometric study; light curve and elements for binary, 51, 193
theory (*see* origin)
tidal evolution; evidence of; U Cephei, 52, 145
Wolf-Rayet star (*see* γ Argus)
Venus; spectrum; systematic shifts of solar lines; variation with zenith distance; explanation; Plate, 53, 380

X-rays

L-series of carbon; identification, 52, 47
source; condensed vacuum spark, 52, 47

Zeeman effect (*see* Spectra)



A 001 217 317 5

